



Date of Surgery: ___/___/___

Surgery Center: _____

Operative Eye: Right Eye Left eye

Pre-Op Instructions Form

- **DO NOT** eat or drink anything 8 hours prior to surgery. You may **ONLY** take your high blood pressure, heart, and thyroid medication with a small sip of water.
- Discontinue all Aspirins, Motrin, Ibuprofen, or pilocarpine eye drops 2 days before your surgery. (Check with your primary care doctor first).
- If you take blood thinners please consult with your primary care doctor prior to surgery.
- Dress comfortably with loose fitted and warm clothing (shirt/blouse that bottoms down in front).
- Expect to be at the surgery center for 3-4 hours.
- **DO NOT** wear make-up, jewelry, or hairpins. Please leave your valuables at home.
- You will **NOT** be allowed to drive home or leave the surgery center unattended. Arrange for a responsible adult to drive you home after surgery and to your post-operative appointment next day.

PRE-OPERATIVE EYE DROP REGIMEN

Please start both your Antibiotic and Anti-inflammatory eye drops 1 day prior your surgery



- **Brand and Generic-Antibiotic:** 1 drop in the operative eye 4 times a day, **1 day prior to surgery.**



- **Brand-Bromsite:** 1 drop in the operative eye 2 times a day, **1 day prior to surgery.**
- **Brand-Prolensa and Ilevro:** 1 drop in the operative eye 1 time a day, **1 day prior to surgery.**